State Water Survey Division
ATMOSPHERIC SCIENCES SECTION
AT THE
UNIVERSITY OF ILLINOIS

illinois institute of Natural Resources

EFFECT OF CONTRAIL CIRRUS ON SURFACE WEATHER CONDITIONS IN THE MIDWEST - PHASE I

by

and

Final Report
to
National Science Foundation

Grant number: NSF AM 78-09568

August 1980

TABLE OF CONTENTS

LIST OF TABLES

LIST OF FIGURES

ABSTRACT

The effect of aircraft contrails on surface climate was assessed by first determining the density of jet traffic across a ten state area of the Upper Midwest, and comparing that distribution with observed changes of various meteorological parameters over the past 50 years, both within and outside of the principal contrail areas. Commercial jet traffic density was found to be greatest in 1979 along a band running from central Ohio westward to central and northern Illinois, with one branch leading westward across northern Missouri, and another branch leading toward the southwest through central Missouri. Another high density region, within the area studied, was eastern Kentucky and Tennessee.

The surface meteorological characteristics were compared for stations within the area of high jet traffic density and those areas less affected by jet traffic, from the middle 1950's until the present. Percent possible sun and number of clear days decreased within the areas of high traffic density from about 1935 to 1950, decreasing more slowly thereafter to the present. The same parameters showed no change or slight increases elsewhere in the study area. The frequency of moderated temperatures, i.e., daily temperature maximum minus daily temperature minimum, increased from the late 1940's with the most pronounced increased found over northern Illinois and Indiana.

In-flight observations from in excess of 16,000 pilot reports suggest that observations with contrails are more frequently also associated with turbulence. In addition and perhaps of more importance to surface climate, aircraft producing contrails typically observe more frequent cloud cover both

below and at the aircraft level. Indeed, only about 1/3 of all contrails can be seen from the surface of the earth due to a natural clouds occurring below the contrails.

There appears to be no difference between the percentage of daytime flights which produce contrails as opposed to those flown during the night. There is a strong relationship however, between contrail areas and pressure features of the middle and upper troposphere, i.e., contrails are much more frequently observed within the area of an upper level trough as opposed to a ridge, and there is a slight preference for contrail areas to be located closest to surface low pressure areas.

Cirriform clouds increased in frequency from 1951 to 1976 according to standard surface meteorological observations. Frequency of clear days decreased over the 26 years at 10 of the 12 first order stations studied. The greatest decline occurred from Chicago to southern Iowa and from Chicago to southern Illinois.

INTRODUCTION

The objectives of the one-year contract were to essentially address two questions: (1) Do contrails measurably alter surface weather conditions in the Midwest? and 2) How well do climatic records from the Upper Midwest indicate the presence of contrails?

To fulfill the objectives, the scope of research included three major thrusts:

(1) Document and analyze jet traffic density over the Upper Midwest from the mid-1950s to the present. If the effect of contrails on climate was to be studied, the location of jet aircraft and contrails must first be identified. This required the collection of a large mass of formatted pilot report information, which included a pilot option to comment on the presence of contrails (either persistent or non-persistent). In addition, since these formatted pilot reports also contained entries regarding several other meteorological parameters, relationships between contrail occurrence and other parameters could also be investigated.

(2) Identify and quantify trends in cloud cover over the Upper Midwest from about 1950 to the present. Researchers have suggested that aircraft contrails have modified natural cloud cover, both forming and spreading into veils of cirrus, and by seeding supercooled high level clouds. This effect, if true, may be preserved in the meteorological observations of cloud cover made routinely at National Weather Service (NWS) offices. Records since about 1950 from NWS first order stations of low, middle and high clouds were analyzed for trends in amount over the last 30 years. The areal distribution of cloud amount (all types) changes relative to jet traffic, and changes of amount of high cloud relative to jet traffic were studied.

(3) Identify changes in the total amount of sky cover, numbers of clear, partly cloudy and overcast days, and changes in "moderated temperature" (decreased diurnal temperature variation) over the Midwest. These parameters were analyzed from the cooperative meteorological network, a much more dense network than the first order stations. The spatial distribution of changes in these parameters was of particular interest, to determine whether observed changes in these potential "response" functions were spatially and temporally related to changes in jet traffic and attendant contrail production.

CHAPTER ONE

SKY COVER, SUNSHINE, AND TEMPERATURE TRENDS FOR 1901-1977

Introduction

This chapter presents results of one phase of the one-year project conducted to investigate the possible extent of influence of contrails on the weather and climate of the Midwest. This particular phase of the study investigated long-term, 77-year records of available data on sky cover, sunshine, and surface temperatures, to examine their temporal variability on an areal basis. The principal goal was to discern any evidence of possible man-made influences on cloud frequencies in the post-1960 period when high flying jet aircraft became more frequent and left many contrails in the upper atmosphere. Examination of the sunshine and temperature data alone could not be considered at all conclusive of a contrail effect, but are useful as material to substantiate whether atmospheric influences from contrails were sizeable and important.

Other chapters of this report delve into possibly more meaningful cloud data, but they are limted to only recent, post-1950 years. Hence, data sets on sky cover conditions and sunshine percentages, as measured at several stations around the Midwest, become the only means for looking backwards beyond the 1950's at the fluctuations in the amount of light being received at the surface and the overlying cloudiness.

This chapter first examines the possible causes for changes in clouds, then briefly describes the cloud climatology of the 10-state study region. This is followed by a section that treats the sky cover and sunshine data (and its quality) and the analyses of these data. The next major section addresses

the results for the study of clear days, and this is followed by section on days with cloudy and partly cloud skies. Sunshine results appear in the next section, then results on surface temperatures, and finally a summary and conclusions section.

Causes of Changes in Clouds

Anyone familiar with the continental climate of the' central United States is aware of the great year-to-year variability in most atmospheric conditions. Indeed, variation on a variety of time scales is the norm of the climate; hence, variations are to be expected, but become major problems when looking for longer term, potentially subtle, trends in weather condition. Any change in the frequency or magnitude of a weather event extending over five or more years in this climate can generally be ascribed to three causes, which may act independently, or in some collective manner, to produce a shift in the climate.

Climatic Variation Due to Natural Causes. The first of these factors are the natural climatic variations due to large scale, hemispheric or global, variations in weather patterns. Unfortunately their causes are not yet well understood. History has shown us that nature has provided a wide range of climates in the Midwest over the past several thousand years. An amazing number of different scales of variations due to natural causes are not only evident but likely. It is also important to recognize that there are at least two types of natural variations that can be expected in cloud frequencies and other weather phenomena. The first might be labeled "the anomalous" type of variation which represent shifts ranging anywhere from five to fifteen years. Such anomalous periods do occur, but the frequency of the weather event

returns to its preanomaly level after the anomalous excursion. Another type of shift is labeled a "climate fluctuation." This often represents a longer term trend ranging from perhaps 50 to 1000 years. Imprinted on both of these are year-to-year variations, but an inspection of their frequencies with some smooth statistical curve fitting, will often reveal a downward or upward type of trend extending over many years. Hence, any systematic change in cloudiness and sunshine noted during the 1901-1977 period must consider that the natural atmospheric shifts could be the cause.

Errors in the Observations. Another reason for apparent systematic changes in cloud cover or sunshine cloud relate to errors or shifts in data collection procedures. The cloud cover data (labeling of each day as clear, partly cloudy, or cloudy) could contain false changes for at least two reasons. First is some form of systematic observer error at a given location, such that the observers are labeling cloudy days as partly cloudy, since the sky observation is somewhat subjective. The other possiblity is that the instructions for record keeping have changed leading to differences in how sky cover is defined. The sunshine records since 1900 have been kept by sunshine recorders, an instrument that inscribes the minutes of sunshine on a chart based on sun falling on the instrument. Errors could come about because of poor instrument calibration, or changes in instrument sensitivity. Similar changes in observation procedures, instruments, or sites could influence temperatures.

Inadvertent Climate Modification. A third reason for climatic shift in the cloud cover frequencies and resulting changes in sunshine and temperature is man's accidental influence on the atmosphere. There appear to be at least three, potentially interactive, ways that man could have systematically been influencing cloud cover since the start of the 20th Century. In the central

United States, the period from 1900 to 1977 has been one of a major shift from an agriculturally oriented economy to a manufacturing oriented economy. In 1900, 12% of the population in the Midwest lived in urban areas, and by 1977 about 70% lived in urban areas. This is also a period of major growth in industrial activity and development of the automobile, particularly since 1940. Hence, there have been major influences on the atmosphere from midwestern cities and industries (Changnon, 1973) .

There appear to be at least three ways in which man's emissions into the atmosphere could have systematically influenced the clouds, sunshine and temperatures. First, are the effects of increased pollutants (aerosols) in the atmosphere. Changnon (1976) has shown how the frequency of smoke and haze days changed dramatically at a series of locations in Illinois with major increases in the number of days beginning in the late 1930's and 1940's. Such increases in man-derived pollutants, generally noted to develop before and during World War II (1935-1945) and remaining high since, have produced additional haze not only in and around cities but now at times over the entire Midwest (Braham, 1977). This haze reduces visibility and restricts one's ability to view clouds. It, of course, also intercepts sunshine and reduces incoming radiation.

A second potential effect of the added pollutants, as has been noted at St. Louis (Grosh 1978) , has been to help produce localized urban increases in cloud cover. A 5 to 10% increase in clouds related to the ability of man-made pollutants to serve as cloud nuclei is well recognized (Landsberg, 1970). Most of these pollutants are urban-related (although automobiles and scattered power plants also add pollutants over a region) and have been realized mostly in low level clouds.

A third inadvertent influence of man on cloudiness, the one that is the subject of this study, is the influence of jet-produced contrails. The potential seems obvious in parts of the Midwest having a notably high frequency of persistent contrails. What is uncertain however is how significant this influence is; that is, whether the contrail-induced cirrus represents an increase in cloud cover of 0.1% , 1%, or 10% or more. In addition to directly producing added cirrus and in inducing more cirrus to form, another influence has been suspected. That is, the ice crystals formed by the contrails could descend into middle level clouds resulting in a) more middle level cloud cover, and/or b) increase in the precipitation.

In essence then, there are several potential causes for anomalous short or long term climate variations in clouds and their influence on the sunshine received at the earth and surface temperatures. Discerning which of these causes is existent and most critical is difficult but the effort is an essential part of this project if the contrail influence is to be isolated.

Climate of Sky Cover and Sunshine in the Midwest

The following sections relating to results of the sky cover and sunshine analysis present the actual numbers on clear days, partly cloudy days, cloudy days, and percent of sunshine for various seasons, years, and stations throughout the 10-state study area (see Tables 2-5) . These numbers will not be repeated here, but a short general discussion is offered. The pattern of sky cover in the study area (Fig. 1) shows that, in general, in all seasons, there are more clear skys (and hence more sunshine received) in the southern and western parts of the study area, as opposed to the northern and eastern. The northeastern part of the study area includes the Great Lakes which produce



Figure 1. Stations and areas used by sky cover and sunshine studies.

sizeable influences on cloud cover, to decrease it in summer and increase it in the colder half year (Changnon, 1968). Current average annual values of sunshine percentages in the north are 48% at Sault Ste. Marie and 51% at Minneapolis, as opposed to 63% at Memphis and 62% at Kansas City. If one considers the annual frequency of cloudy days, the current average at Sault Ste. Marie is 213 days with 174 days at Minneapolis. Comparable average values to the south are 151 cloudy days at Kansas City and 153 days at Memphis..

There are also systematic differences across the region in the seasonal frequencies of sky cover. For example, in the South Area (see Fig. 1), the leading season of clear days, on the average, is fall followed by summer, spring and winter. In the South Central Area, clear day frequencies in fall are equalled by those in summer, followed in order by the frequencies in spring and winter. As one moves farther north into the North Central Area, clear day frequencies become greatest in summer with fall values ranked second, then spring and winter. In the northernmost part of the study area, summer clear days, on the average, rank first, followed by the number in spring, with fall frequencies ranking third and winter last.

If one examines the average seasonal frequencies of clear days on a west to east basis, one finds that the fall frequency leads in the west followed by summer, spring, and winter values. However, in the eastern half of the study area (Fig. 1), summer is the leading season with clear days followed by fall, spring, and winter.

In general, in most parts of the study area, the fall and summer are the seasons with the greatest frequencies of clear skies and hence more sunshine. Winter ranks last among the four seasons in the amount of sunshine and number of clear days, and obviously ranks first amongst the four seasons in the number of cloudy days.

Cloud and Sunshine Data and Related Analyses

The stations for which data on sky cover and sunshine were available for all or most of the 1901-1977 period are shown in Figure 1. Except for a few stations such as South Bend, Indiana, most all stations had complete records of sky cover for this 77-year period. However, records of sunshine at most stations began in the 1910-1915 period, and very few stations had available sunshine records prior to 1910. The periods of record are indicated on Table 1.

Great care was also made in the study of the historical records to consider any possible observational or instrumental errors. During the 1901-1977 period, there were no changes in instructions to Weather Bureau (now National Weather Service) observers about how to list the sky cover as being cloudy, partly cloudy, or clear. Since observations were made by a large number of persons over the 77-year period, it is doubtful that any systematic observer biases existed to produce long term trends. Hence observer error is discounted as a major factor causing any trends discerned.

The potential of faulty equipment or poor calibrations of the sunshine recorders, and hence systematic time biases cannot be discounted. Instruments were maintained, as near as one can learn, reasonably well and instrument problems or changes should not be a reason for any long-term (greater than 5 to 10 years) trends in sunshine data. Wahl's (1968) study of data from several midwestern stations of both the mid 19th Century and mid 20th Century suggets near equal precision of temperature from both periods.

One of the "checking procedures" used in this entire study to verify the potential of "errors" in the sunshine or sky cover data was to inter-compare them. Their results (trends) should agree, and yet one set of data (sky cover) is by human observations and other is based on instruments. As will be

noted in the results, there is general agreement in the trends of sky cover and sunshine, suggesting that the observer error or equipment problems, as causes for trends, was negligible. However, one important change at 34 of 36 stations (Cairo, Illinois, and Marquette, Michigan excluded) was a relocation of the station site from the city (or town) central business district to the rural airport. As shown in Table 1, these largely occurred in the 1940's but began in the mid 1930's and continued into the early 1950's. These shifts involved relocation, at least at the largest cities, to locales where less urban pollution existed (or would exist). Hence, the site change represented a possible improvement in visibility, and the potential for less cloud cover and a potential for more sunshine.

The various analyses were based on daily values of sky cover and percent of possible sunshine. The daily sky cover values of clear (0 to 0.3 sky cover), partly cloudy (0.4 to 0.7 sky cover), and cloudy (0.8 to 1.0 sky cover) are values derived from hourly observations made at the U.S. Weather Bureau first-order stations scattered throughout the area. These hourly observations were combined into daily values. Monthly frequncies of clear, partly cloudy, and cloudy days were obtained in this study either from published records or original records of the first-order stations. Once compiled and totaled, the values were entered into punch cards for subsequent analysis.

The daily sunshine recorder data were used in a similar fashion. Daily sunshine values are recorded both as to actual time and as percent of possible sunshine for the day. We chose to analyze the percent of possbile sunshine which accounted for changes in length of day during the season. Daily values of possible percent of sunshine were combined into monthly means. These

Table 1. Periods of records and dates of shift in the location of First Order Stations from inner city sites to airport sites, during 1901-1977.

	City to Rural airport site	Period of cover dat	Period of sun¬ shine data
Memphis, TN	Sept. 1950	C	1902-77
Nashville, TN	June 1941	C	1919-77
Chattanooga, TN	June 1940	C	1919-77
Sault St. Marie, MI	July 1951	C	1922-77
Marquette, MI	Always in city	C	1905-77
Grand Rapids, MI	Aug. 1950	C	1922-77
Detroit, MI	July 1942	C	1922-77
Alpena, MI	Aug. 1959	C	1922-77
Ft. Wayne, IN	Aug. 1939	1902-	1912-77
South Bend, IN	Feb. 1933	1914-	none
Indianapolis, IN	May 1941	C	1916-77
Evansville, IN	Aug. 1940	C	C
Milwaukee, WI	Aug. 1940	C	1916-77
Madison, WI	Sept. 1939	C	1916-77
La Crosse, WI	Dec. 1950	C	none
Green Bay, WI	Aug. 1949	C	1903-77
Little Rock, AR	Dec. 1938	C	1916-77
Ft. Smith, AR	Sept. 1945	C	1916-77
St. Louis, MO	July 1958	C	1916-77
Springfield, MO	Aug. 1940	C	C
Kansas City, MO	Jan. 1934	C	1916-77
Columbia, MO	Jan. 1936	C	1916-77
Louisville, KY	June 1945	C	1916-77
Lexington, KY	July 1944	C	none
St. Cloud, MN	Feb. 1936	1916-	none
Rochester, MN	Nov. 1943	C	none
Minneapolis, MN	Oct. 1937	C	1916-77
Duluth, MN	March 1950	C	1911-77
Dubuque, IA	June 1951	C	none
Burlington, IA	Oct. 1940	1914-	none
Des Moines, IA	Oct. 1950	C	C
Moline, IL	Dec. 1939	C	C
Springfield, IL	Jan. 1948	C	C
Chicago, IL	June 1942	C	C
Cairo, IL	Not moved	C	1920-77
Peoria, IL	May 1943	C	C

1/ = complete for 1901-77

monthly mean values of percent of possible sunshine were calculated from historical records of the first-order stations, punched onto IBM cards, and subsequently used to develop seasonal and annual values.

Much of the analysis was based on seasonal and annual values of sky cover and percent of sunshine. The seasons chosen for analysis of the sky cover and sunshine data were spring (March-May), summer (June-August), fall (September-November), and winter (December-February). For the sky cover data and the sunshine data, the seasonal and annual values were used to develop moving averages of 2-, 5-, 10-, and 20-year periods. Graphs of these moving averages were plotted to search for the most informative means for portraying the results and removal of small scale fluctuations. In most cases, the 10-year totals were chosen to study the trends in sky cover and sunshine.

One phase of the analysis included inspecting the data and trends of the 36 individual stations in the study region (Fig. 1). Basically, various descriptions of the temporal behavior of the 10-year values of the individual stations, when plotted on base maps, suggested the existence of various regions having similar temporal characteristics, as defined by the stations within them. To further investigate areal differences on a more systematic bases, the study region was divided into six areas, as shown in Figure 1. Four equal sized (north-to-south) areas were chosen to allow for a target-control approach in the comparative analysis. Available evidence on persistent contrail frequencies across the Midwest (see Chapter 3), and on commerical jet aircraft traffic across the study region (see Fig. 2), both show that the "central" portion of this study region is the one with the greatest likely contrail influence. This central area is an east-west area bounded on the north by southern Wisconsin and on the south by southern Illinois (and also includes northern Missouri, Iowa, the northern 2/3's of Indiana, and southern Michigan). This general hypothesis of effect in the



Figure 2. The daily frequency of commercial jet traffic in 1979, based on number of flights per 12, 380 km² area.

central areas was tested by discerning whether the frequency of sky cover conditions and sunshine in the two central area sections (labeled North Central and South Central in Fig. 1) differed from those in the northern and southern areas where less jet aircraft existed along with lesser urban industrial development. In the north and south areas there is less likely man-made effect from other causes (fewer big cities, lesser population), and any sky cover and sunshine changes there are thought to be due mostly to natural variations in climate. Hence, the areal comparisons became a major means of evaluating man-made influences in the study region.

The daily frequency of commerical jet aircraft flights across the study area in 1979 is shown in Figure 2. This was determined by measuring each of the flight tracks on a Lambert Conformal Projection, and counting these based on a grid 60 x 60 nautical miles squares. The flight data came from the August 1979 issue of the Official Airline Guide. Flights were counted only if they exceeded 200 nautical miles in length (shorter flights typically do not reach contrail producing levels). The ending or starting 50 nautical miles of each flight also were not counted since this is typically a distance of being below contrail levels (takeoff and descent for landing). The pattern of Fig. 2 clearly shows the predominance of flights in the two central areas, and particularly in the South Central Area. The averages and extremes of the grid squares in the four study areas are:

	Average	Maximum	Minimum
North Area	41	105	10
North Central	224	700	22
South Central	345	504	116
South Area*	211	508	130

Although the South Area is used as a "central area" in these analysis, the high frequency of flights in its eastern portions indicates possible contrail influences there. Also, the upper tier of squares in the North Central Area have low values suggesting a potentially lesser influence than one might expect for the South Central Area. Inspection of Figure 2 and grid areas > 300 flights per day shows the east-west oriented flyway that is bounded on the north by northern Illinois and extreme lower Michigan and bounded on the south by a line through St. Louis.

Clear Day Results

Annual results. At all but two of these 36 stations, general downward trends in clear day frequencies were found during the 1901-77 period. Only Cairo, Illinois and Green Bay, Wisconsin (Fig. 1) had essentially flat unchanging trends in clear days. Curves of the average annual frequencies of clear days of the four areas are shown in Figure 3. Essentially, frequencies were highest in the 1901-1940 period; followed by sharp declines during the 1940's; followed by a flatter trends since 1950. Notable on these time curves are the regional climatological differences with higher frequencies in the south and the least number of clear days in the north. The concern over the effects of the city-to-rural site changes of the stations seems of little consequence in the 1940's. The visibility improved at a time when clear skies decreased greatly indicating little effect on the sky cover observations.

Also notable on the graph (Fig. 3) are other temporal features. For example, the high value of 1931-40 in the South Central Area (with decreases existed in other areas) is a reflection of the severity of the drought of the



Figure 3. Area near 10-year values in clear days.

1930's in that area, as opposed to other parts of the region where it was less severe. Quite likely, there was less industrial development in that southern area in the 1930's than elsewhere.

Following the major decrease to the low number of clear days in the 1941-50 period (which was also a notably wet decade in the Midwest), slight increases in the annual number of clear days occurred for the following decade, 1951-60. This was followed by decreases having different rates (by area) since 1960. The greatest rates of recent decrease are found in the two "target" areas, the North Central and South Central Areas. The declines are much sharper during the last 10 years, 1968-77.

In one sense, the shapes of the 1901-77 frequencies of clear days (Fig. 3) fall into three types. The first of these is a general smooth downward trend from 1901 to 1977 shown by the North Central Area. Another type is a downward trend showing the lowest values of the study period to be in 1941-50. It is found both in the North and South Areas. The third type has several trends and two minima, with a major low in 1941-50, and then a subsequent greater low in the 1970's. It is reflected in the South Central Area curve. Differences between the curves shown on Figure 3 are very important, indicating different climatic regimes or influences for clear days in the two central areas. Their clear day values and those to the north and south show differences after 1940 (greater decreases in the two central areas) suggesting man-made influences in these two central areas. Both central areas are more industrialized, have greater population centers, and have more contrail producing aircraft traffic (Fig. 2).

The regionality of the annual clear day results is further reflected in Figure 4. Here, the characteristics of the trends in annual clear days for the 1901-77 period are shown, as based on the characteristics found in the 36



Figure 4. Characteristics of trends in annual clear day frequencies for 1901-1977 period.

individual first-order station (10-year) curves. Through the central section, a continuous downtrend is evident and it occurs where the greatest air traffic exists. To the north and to the southeast of that area, the station curves showed a downtrend to a minumum in the 1970's but with a major secondary minimum in the 1941-50 period. To the south, there is a general downtrend but with the major minimum occurring in 1941-50. Hence, Figures 3 and 4 reveal that there have been two major minima found at most stations in the study region, one coming in the 1941-50 period and one in the 1968-77 period.

Another means for displaying the regionality of the clear day frequencies is portrayed in Figure 5. Here, the year ending the lowest 10-year clear day values at the 36 stations were plotted. Regions were developed from these ending year values. One notes that a large portion of the central and eastern section of the region had its lowest 10-year frequency of clear days ending during the 1974-1977 period. However, there are other areas in the north and south where the lowest year came in the 1940's and up through 1951. One area in the west had its lowest 10-year value of clear days in 1963-65.

Table 2 presents the annual percentage changes between two decades, the earliest (1901-10) and latest decades (1968-77) for the 6 areas. These reveal that the greatest percentage decrease (32%) occurred in the South Central Area, but there was no difference on a west-east basis. The average 10-state decrease was 26%.

Comparison in the lower part of Table 2 of the annual "predicted" values reveals that the decreases between the early and late decades were 9 to 11% greater than would be expected, or predicted, on a "climatological trend basis." The predicted values shown in Table 2 were derived from a linear interpolation of the "control area" values in the north and south and are considered to be best estimates of changes due to natural variations.



Figure 5. Areas based on ending years of lowest 10-year values of clear days (annual) for 1901-1977 period.

Table 2. Change in 10-year values of clear days from 1901-10 (early) to 1968-77 (late).

Areas	Early-late differences X 100 Early Value Spring	Summer	Fall	Winter	Annual
North	-27	-23	-30	-29	-18
North Central	-23	-23	-35	-33	-29
South Central	-28	-32	-42	-29	-32
South	-13	-19	-33	-14	-22
West	-27	-28	-35	-24	-26
East	-20	-27	-35	-25	-26
Total Area	-23	-27	-35	-24	-26

Area	Summer			Fall			Annual		
	p(1)	A(2)	C(3)	p	A	C	P	A	
NC	-21	-23	+ 2	-31	-35	+ 4	-20	-29	+ 9
SC	-20	-32	+12	-32	-42	+10	-21	-32	+11

(1) p = predicted value based on a linear interpolation between the north and south area.

(2) A = Actual area value.

(3) C = Change or difference between actual and predicted values.

Seasonal results. Area average values for 1901-77 (such as displayed in Fig. 3) were developed for each season. Basically, these showed general downtrends in all seasons.

Inspection of the areal seasonal values presented in Table 2 reveals that the least percentage change (betwen the early and late decades) occurred in spring and the greatest in fall. East-west seasonal differences are small (except in spring), but the differences among the north-to-south areas are considerable. The least changes occurred in the South Area in most seasons. The greatest percentage decreases occurred in the South Central Area in all but the winter when the greatest decrease occurred in the North Central, or other "target" area. The total area average values show that the fall (-35%) and summer (-27%) were the seasons of greatest decreases in clear days.

Shown at the bottom of Table 2 are the differences for these two seasons and for the North Central Area and South Central Area, the two target areas. The difference between actual values and the predicted values (based on climatological interpolation) revealed the "change value" believed indicative of man-made influences. These show that in the South Central Area there was an anomalous 12% decrease in summer and a 10% decrease in fall, both due presumably to man-made influences.

Since the greatest seasonal decreases, both in total percentages and those expressing potential man-made influences, were found in the South Central Area, the 10-year values of each season in that area are presented in Figure 6. These are the averages based on the 11 stations located in this area (Fig. 1). One notes a generally flat trend for 1901-1940, although winter and spring values begin to decrease after 1930. All four seasons show a marked decrease in clear days in 1941-50, followed by generally flat trends thereafter. However, the fall and summer seasons show definite decreases



Figure 6. 10-year area mean values of clear days in South Central Area.

after 1960 in the number of clear days, but the winter and spring frequencies remained constant after 1960. Since jet aircraft traffic in the area increased sharply in the early 1960's, the values shown for 1961-70 and for 1968-77 are potentially related to contrail effects and might be reflected in the number of seasonal clear days. The summer decrease in clear days over this South Central Area went from 28 days in 1951-60 to 25 days in 1968-77 (a 3-day decrease). The fall season change went from 35 days in 1951-60 to 24 days in 1968-77. Both seasons exhibit their 77-year minima in the last 10 years.

In general, the seasonal curves of Figure 6 strongly suggest a man-made influence leading to a decrease in clear skies beginning in 1941 and likely related to the great expansion of industrial activity related to WWII and which continued thereafter. The more recent decreases shown since 1960 may be a mixture of effects of the greater urban growth and the jet contrail influences.

Summer season. Analysis of the individual stations and their summer frequencies of clear days was revealing. In Figure 7, the 10-year values of three stations selected from the north, southeast, and southwest parts of the study region are presented. Inspection of these trends reveal interesting differences; for example, the Minneapolis frequency of summer clear days fluctuates but has no discernible up or down trend during the 77-year period. However, both the Columbia and Evansville curves exhibit down trends with a much greater down trend at Columbia, ranging from 39 clear days per summer in 1901-10 to 25 clear days per summer in 1968-77. Evansville, which has a minimum in recent years, had its greatest minimum in 1941-50.

The characteristics of the curves of summer clear days of the 36 stations were plotted to develop a regional map. The general characteristics of the trends in summer clear days frequencies is shown in Figure 8. Major down



Figure 7. 10-year values of clear days in summer and fall at selected stations.



Figure 8. General characteristics of trends in summer clear day frequencies for 1901-1977 period.

trends (as revealed by the Columbia curve in Fig. 7) were found along an east-west corridor through the central section. Lesser downtrends were noted in the northwest and south central districts, with essentially flat trends in the northern, southwestern, and southeastern sections. The pattern of Figure 8 is suggestive of man-made influences, with the greatest downtrends occurring where the major jet air traffic exists (Fig. 2), as well as in the major industrial section of the 10-state region.

Fall season. Individual curves of the 36 stations for their frequencies of clear days in the fall season revealed downward trends everywhere. Examples of the curves of three stations are shown in Figure 8. The values at Minneapolis (North Area) indicate a generally flat trend through 1960 followed by a decrease in recent years. The Little Rock (South) and Columbia (South Central) curves also show recent decreases, but with a much greater downtrend in the recent 20 years at Columbia. In general, the trends in the central areas of the study region showed considerably decreases, particularly since 1960, in frequencies of clear days.

Winter season. The South Area stations showed decreases in clear days to around 1950, followed by up trends to the present. In the central area most stations revealed a maximum of clear days in 1921-30, followed by decreases during the decades thereafter. Five stations had their lowest frequency of winter clear days in 1968-77 including Peoria, Columbia, Des Moines, Ft. Wayne and South Bend. The northern stations indicated flat trends in winter clear days. There appears to be little evidence, other than at a few central stations, of any man-made influence on clear days in the winter season.

Spring season. Stations in the South Area had generally downward trends in clear days in spring until 1950, followed by uptrends to present. They were very similar to winter clear day trends in this same area. In the

central section, spring frequencies of clear days generally decreased after 1920 with either a minimum occurring in 1941-50 or in 1951-60. Only a few stations (Peoria, Burlington, Louisville, and Columbia) had their lowest values in the 1970's. In the north, spring values showed decreases and long-term downward trends. Several stations reached their lowest values in the 1970's. In general, the spring results of clear day frequencies were not strongly suggestive of localized influences throughout the North Central or South Central target areas.

Cloudy Day Results

Annual values. The 10-year values of cloudy days for the four areas are shown in Figure 9. All four areas have notable upward trends from 1901 to 1977. The total area average increase (Table 3), or difference between the earliest and latest decades is 41%. In general, the cloudy day curves of Figure 9 exhibit more smooth and notable trends than do the comparable annual clear days curves of Figure 3. The cloudy day values are more indicative of a general, long-term climatic change than are the clear-day frequencies because part of the shifts are in partly cloudy conditions as well as clear conditions. Further comparison of the curves in Figures 3 and 9, such as those for the North Area, reveal a good inverse relationship, particularly up through 1960. It is important to also note that both the "target areas," the North Central and South Central Areas, exhibit sizeable increases in cloudy days in the last decade, 1968-77, particularly evident in comparison with the lesser up trends for this North and South Areas for this period.

Inspection of the annual values in Table 3 also reveals that it has become relatively cloudier in the south, central, and west areas with a lesser increase in cloudy days in the eastern and northern areas of the study



Figure 9. Area mean 10-year average annual values of cloudy days.

Area	Early-Late Differences y_{100} / Early Value				
	Spring	Summer	Fall	Winter	Annual
North	+21	27	21	22	22
North Central	+38	52	52	33	43
South Central	+41	82	91	3„	53
South	+25	69	86	21	44
West	+41	58	65	32	46
East	+28	52	42	24	33
Total Study Area	+34	56	6„	27	41



Figure 10. Seasonal area mean frequencies of cloud days in South Central Area.

region. This is, the areas that in the early part of the 20th Century were the clear, less cloudy areas have become cloudier faster than the areas which were, on the average, cloudiest (north and east).

This suggests, as do the later results of greater increased cloudiness in summer and fall seasons, that the additional cloudiness has been occurring in these areas and seasons which normally are most clear. This indicates that whatever the causes, the influences are more obvious during periods and places which in the past have been more clear. This is supportive, in general, of the type of influence that added high clouds could be expected to produce. Additions of high clouds in areas and times that are already relatively cloudy would be less noticeable than in relatively clear areas.

Seasonal results. The trends in cloudy day seasonal values at most stations in the 10-state region were generally flat during the 1901-40 period. They then began to increase, particularly in the South and South Central Areas. Please note these features in the seasonal curves (Fig. 10). As shown in Table 3, the greatest changes percentagewise in cloudy days from the early decade (1901-10) to the latest decade (1968-77) occurred in the fall season. The total area average increase is 60%, closely followed by summer with 56%, with much lesser increases in spring and winter. The percentage changes shown in Table 3 for cloudy days are much greater than those shown for clear days (Table 2), except in the North Area where the clear day percentage changes exceeded those for cloudy days in spring, fall, and winter.

As with clear days, the percentage changes in cloudy days were greatest in the South Central Area in all seasons except winter when in the North Central Area was greatest. There are also west-east differences shown in all four seasons with the greatest increases found in the western region in all four

seasons. It is important to note that the South Central percentages approach 100% for summer and fall. Thus, there have been notable increases in cloudy days in this area of greatest jet aircraft traffic.

The North and South Areas were considered to be control regions and either non-affected by contrails, or less affected by contrails than the atmosphere over the North Central and South Central Areas. Thus, their values were used as controls to estimate the likely "natural variability values" for the North Central and South Central Areas. The differences calculated between these climatologically predicted (interpolated) values and the actual values for the North Central and South Central Areas are an indication of possible man-made influences.

Table 4 shows the changes in the two central areas for all four seasons, based on the comparison of the predicted and actual values. The changes indicate anomalous increases in the recent decade in all four seasons with about 10% being representative of the regional value of increases in the spring and winter. However, in the fall and summer seasons, the South Central Area had about 26 to 27% anomalous increases in cloudy days with about 10% in the North Central Area. The annual values in Table 4 suggest a 14 to 17% anomalous increase in cloudy days over that expected. Hence, results indicate the greatest potential influence has occurred in the South Central Area.

As a result the South Central Area seasonal curves are presented in Figure 10. These indicate a generally flat trend until 1930, or 1940, followed by increases in cloudy days, generally sizeable for the 1941-50 period. The fall and summer season curves also show substantial increases after 1960. These are not found in the spring and winter seasons. The shift from 1960 to 1977 in the cloudy day values in fall is from 31 days to 44 days (up by 13 days), and in the summer the shift is from 29 days to 32 days.

Table 4. Comparison of predicted and actual percentage changes from 1901-10 to 1968-77 in cloudy days.

		Spring		Summer			Fall			Winter			Annual		
Area	p(i)	A[2]	C[3]	P	A	C	P	A	C	P	A	C	P	A	C
NC	26	38	+12	41	52	+11	43	52	+9	22	33	+11	29	43	+14
SC	30	41	+11	55	82	+27	65	91	+26	21	30	+9	36	53	+17

[1]P = Predicted value based on lindear interpolation between control values, the North and South areas.

[2]A = Actual area value.

[3]C = Change, or differences, between actual and predicted value.

Partly Cloudy Days Results

Comparison of the cloudy and clear day results has shown great similarity. Much of the increase in cloudy day frequencies was reflected in the decrease in clear days. Inherently this means that frequencies of days with partly cloudy skies were not changed greatly, and hence their results are not presented in great detail.

The summer season trends in partly cloudy days of the 36 stations define three areas. The area south of a line from Springfield, Mo. to Lexington, Ky., had essentially flat trends for 1901-77 in the number of partly cloudy days. The stations in the area north of a line from Burlington to Alpena also had flat trends. The remaining central area had down trends in their summer frequencies of partly cloudy days, generally decreasing by 10 to 15% from 1910 to 1977, with the lowest values in 1968-77.

The fall season trends in partly cloudy days were similar to those in summer. Moderate down trends of 10 to 20% were evident at most stations in the South Central Area. Elsewhere the 1901-77 values varied but did not reflect any down or up trend. The winter and spring frequencies of partly cloudy days showed no trends.

The down trends in partly cloudy days, reaching a low in the 1968-77 period during summer and fall, and largely within the South Central Area do suggest a contrail related influence. The most important finding is that shifts to cloudier conditions in the Midwest were largely associated with a decrease in clear days, with lesser change in partly cloudy days. This helps indicate the shift was sizeable.

Sunshine Results

Annual Results. The percentage of possible sunshine values were analyzed on an annual and seasonal basis. Additional cloudiness and loss of clear days, as shown in the previous sections, should be reflected in a decrease in sunshine, on an annual basis and all seasons, but particularly in fall and summer. The general characteristics of the trends of the annual percent of sunshine values for 1901 (or whenever records started in the 1901-22 period) to 1977 are shown by the pattern in Figure 11. The station trends were found as having a general down, up, or flat trend. Importantly, one notes the general downward trend throughout the central section where clear day trends were downward (see Fig. 4). To the north and south of this central area (the major jet flyway as shown in Figure 2, and urban-industrial section of the study region) one finds either flat or upward trends in the sunshine data.

The years representing the year ending the lowest 10-year percent of sunshine value in the 20th Century are shown in Figure 12. This pattern strengthens the trend results showing the lowest values have come in the late 1970's in the central section and largely where jet traffic is greatest. In other areas to the north and south of the central section, the major minima occurred in a variety of other years, but mainly in the late 1940's as with clear days. In general, the sunshine results and clear days results are remarkably similar indicating equal quality of both sets of data.

The changes in the annual 10-year percent of sunshine values, between 1921-30 (early) and 1968-77 (late) were quantified, as shown in Table 5. The annual values, although much less percentagewise than those found for clear days or cloudy days, do show greater decreases in the central target areas than those in the north and south control areas. Comparison of the



Figure 11. General characteristics of trends of annual percent of possible sunshine values for 1911-1977.



Figure 12. Ending year of lowest 10-year values of annual percent of possible sunshine in 1901-1977 period.

Table 5. Changes in 10-year average percent of sunshine values from 1921-30 (early) to 1968-77 (late).

$$\frac{\text{Early-Late}}{\text{Early Value}} \times 100$$

Area	Summer	Fall	Annual
North	+2	0	-2
North Central	-3	- 9	-4
South Central	-4	-10	-6
South	-1	- 6	-2

		Summer		Fall			Annual		
	(1)	A(2)	C(3)	P	A	C	P	A	C
North Central	+1	-3	-4	-2	- 9	-7	-2	-7	-2
South Central	0	-4	-4	-4	-10	-6	-2	-5	-4

(l) = Predicted value based on linear interpolation between central values, the north and south area.'

(2)A = Actual area value.

(3)C = Change.

"predicted" and actual annual values in these two "target areas," shown in the lower part of the Table 5, indicates a change in sunshine of 2% in the North Central Area and a 4% decline in the South Central Area. The shift in the annual sunshine values is further illustrated by the area curves shown in Figure 13. They all show a decrease from 1921 through 1950, with a notably sharp decrease in 1941-50, as found in the sky cover conditions. Furthermore, the North Central and South Central Areas show reductions in 1968-77 that are greater, however, than in 1941-50. The other, control areas have their lowest sunshine values in 1941-50.

Summer. Inspection of the area summer curves for 1921-77 in Figure 13 reveals most stations had down trends in the summer percent of possible sunshine. The curves of the individual stations were analyzed for their trends and a map based on their trends is shown in Figure 14. Again, the central area had down trends, with essentially flat or up trends found to the north and south. As shown in Table 5, the downward trends of 3 to 4% in the North Central and South Central Areas were estimated as being 4% more than climatological prediction suggested, reflecting potential man-made influences.

Fall. Downward trends in the percent of sunshine in the fall season are noted in most areas for the 1961-77 period (Fig. 13). Only the North Area has no downward trend. Station trend values were used to develop Figure 15 showing the patterns of general trends in fall sunshine for 1921-77. Here, again we note the downward trend through the central area with flat trends to the north and to the southwest. As shown in Table 5, the net effect of the fall sunshine trends is to produce an estimated 6 to 7% decrease in sunshine in the North and South Central Areas considered due to man-made influences. The sunshine decrease in the last 17 years and in the period of jet air traffic -suggests a contrail-related influence on sunshine in the fall.



Figure 13. 10-year values of percent of possible sunshine in 4 areas.



Figure 14. General trends in summer sunshine percentages from 1911 to 1977.



Figure 15. General trends in fall sunshine from 1911 to 1977.

Temperature Results

Much has been presented in the last 25 years about man's impact on weather and climate. It has been clearly demonstrated that man has altered his urban climate so as to significantly change the surface temperature and most other local weather features (Changnon, 197'3) . The fluctuations in global temperatures over the last 90 years have been partially ascribed to man's activities and principally to the release of pollutants to the atmosphere (Mitchell, 1971) . The recent increase in moderated (less extreme) temperatures in parts of the Midwest may also reflect a climatic change that appears to be partially attributable to various man-made conditions.

There are two commonly cited hypotheses for how man's pollutants may affect surface temperatures to produce either warming or cooling. First is a belief that the particulates in the atmosphere absorb and reflect incoming visible solar radiation leading to a reduction in solar energy received at the earth's surface, thus resulting in cooling. However, Mitchell (1971) has shown that man-made aerosols in the troposphere may serve to increase surface temperatures in certain areas, whereas aerosols in the stratosphere lead to cooling. Secondly, direct thermal pollution and gases in the atmosphere, such as CO2 which is opaque to long-wave radiation (re-radiated from the earth), can produce a "greenhouse" effect leading to an increase in surface temperatures. Bryson (1972) claimed that both inadvertent cooling and inadvertent warming conditions have existed in the 20th Century helping to produce initially about 1.0 C warming (1900-1945) and a subsequent 0.4 C cooling since the 1940's.

An interesting possibility relating to increased atmospheric pollutants, both particulate and gaseous, concerns their potential for moderating extremes of temperature. If the incoming solar energy is reduced by particulates during the daytime, the period of maximum heating, then the maximum temperature values would be reduced. If gases are sufficient to decrease the amount of outgoing radiation, particularly at night when minimum temperatures normally occur, then an increase in temperatures might be expected. Certainly, other man-made factors such as increased cloudiness would have a similar "thermostatic" effect since clouds reflect solar radiation and also decrease outgoing terrestrial radiation. In fact, some of the increased cloudiness, at least in recent years, has apparently resulted a) from contrails produced by high altitude jet aircraft, which have been estimated to have increased high-level cirrus clouds by 5 to 10% (Bryson et al., 1970), and b) from the added man-made particulates which have furnished more cloud condensation nuclei and ice nuclei to aid low and middle level cloud development.

These possibilities for man-induced moderation of temperatures were investigated by use of 70-year sets of records from several relatively undisturbed weather stations from small midwestern towns. Since contrail influences are believed to produce only subtle temperature changes, stations were selected that had long undisturbed records and were without any major urban influences on them. Several of the stations selected were identified by the National Weather Service as national bench-mark stations, i.e., high quality stations. Other stations were carefully selected for analysis by knowledge of their quality and lack of station relocations. These form N-S and E-W lines across the study region, and their locations are shown in Figure 16.



Figure 16. Decadal frequencies of months with moderated temperatures.

Months having both 1) below-normal average maximum monthly temperatures, and 2) above-normal average minimum monthly temperatures were selected to indicate a period of reasonably prolonged moderation of temperature. The monthly temperature data for the 1901-1977 period were examined to identify those months with moderated temperatures.

The seasonal frequencies of these moderated months in 10-year periods are displayed in Table 6. They occurred most frequently in summer (June-August) and fall (September-November). The annual average, based on the 70-year values in Table 6, ranges from 11 to 14 months of moderated temperature per decade about 10 percent of the time). If the departures of the maximum and minimum temperatures were independent events, the probability of a moderated month would be 0.25, or they would occur 25 percent of the time instead of 10 percent. Thus, the incidence of moderated temperatures is not a common event.

Comparison of the decade totals in Figure 16, based on stations aligned along north-south and west-east lines, reveals that stations in the central area exhibited sharp increases in months with moderated temperatures during 19 61-70. Stations to the south and north did not. Stations along the east-west line where contrail production is greatest (Fig. 2) all showed a notable increase in moderated months in 1961-70, the period when jet aircraft traffic significantly increased. Comparison of the temperature frequency curves in Figure 16 with those for cloudy days (Fig. 9) reveals general agreement. Since the increased cloudiness in the 1961-77 period may likely be a result of the contrail induced cirrus in the central area, it appears the effect has at least partially caused a moderation of temperatures.

Further study involved comparison of the Urbana temperatures with the sky cover conditions (frequencies of cloudy days) at Peoria. Peoria is located 110 km (75 miles) northwest of Urbana and its cloud cover values should be

Table 6. Seasonal frequency of months with moderated temperatures (max < normal with min > normal) in 1901-1970.

	Average number of months per 10-years				
	Sp	Su	F		Annual
Chatham, Mich.*	4.0	3.1	3.1	2.9	13.1
Urbana,	2.4	4.4	4.1	1.8	12.7
Brunswick, Mo.*	1.6	3.4	4.6	2.1	11.7
Lewisburg, Tenn. *	2.6	5.3	4.1	2.4	14.4
Dixon,	3.0	3.6	4.4	3.0	14.0
Aurora, I11.	3.1	3.0	3.6	1.7	11.4
La Harpe,	2.3	4.6	4.0	2.1	13.0
Mt. Vernon,	2.5	4.9	4.8	2.3	14.2
New Burnside,	2.5	4.4	3.0	2.3	12.2
Winamac, Ind. (1910-70)	2.4	4.5	5.1	2.2	14.2
Watertown, Wis.	1.6	3.8	3.7	1.5	10.6

*Bench-mark Stations.

representative of those at Urbana. Comparison of the cloud conditions and the moderated temperature frequencies for 1901-73 (Fig. 17) reveals that the increase in moderated temperatures at Urbana is partially related to a changing frequency in cloudy sky conditions. The correlation coefficient of the two sets of annual values is +0.1 indicating that the cloudiness explains about 25 percent of the variance in the 75-year record of frequency of moderated temperatures. However, of importance is the fact that the highest values of both conditions occurred together and in the last few years (Fig. 17). The spectacular increase in cloudiness since the 1940's is a matter of considerable interest.

Examination of whether the moderated temperature increases at Urbana relate to particulate and gaseous pollutants (and not clouds) was limited to use of the only long-term records potentially indicative of such pollutants, the frequency of days with haze and smoke at Peoria. These are only a gross measure of the visible pollutants but were examined realizing they may be an index representative of regional pollution influencing Urbana. The smoke-haze curve in Figure 17 shows an increase in the visible atmospheric pollutants during the 1930's, and then again in more recent years. The correlation between the annual values of smoke-haze days and the frequency of moderated temperature months was 0.32. Although the Peoria smoke-haze frequencies are only an index of pollution, and may not represent conditions at Urbana, a weak relationship does exist between their frequencies, suggesting that man-made pollutants may have caused some of the recent doubling of months with moderated tempertures at Urbana.

The peak of smoke-haze conditions in the 1931-40 period (Fig. 17) is not entirely relatable to atmospheric pollutants from combustion, particularly since industrial activity in that period was considerably less than in



Figure 17. 5-year values of moderated temperatures at Urbana, and cloudy days and smoke-haze days at Peoria.

subsequent decades. However, the 1931-1940 period had record droughts in Illinois and the central United States. These droughts resulted in enormous dust storms creating large volumes of dust particles in the atmosphere which also cloud serve to reflect incoming solar energy and possibly to inhibit long-wave radiation. A means of further investigating this dust explanation for more moderated temperatures in both the 1930's and 1960's is to look at dust "indicator" data for Peoria (dust storms are defined on a daily basis when dust limited visibility during the day to 1/2 mile or less) . The number of dust storm per decade since 1905 were 0 in 1911-20, 0 in 1921-1930, 25 in 1931-40, 7 in 1941-1950, 0 in 1951-1960, and 1 in 1961-1970. The singular peak in dust storms activity in the 1930's may help explain the peaks during that period in the frequency of moderated tempertures (Fig. 17), and in the smoke and haze days, since haze is closely related to dustiness.

The added cloudiness and the increase in visible pollutants, both partially man-made, together explain some 45 percent of the fluctuations in the moderated temperature frequencies since 1900. However, these two factors both relate well to the recent (post 1960) large increase in moderated temperatures. This suggests that recent mad-made activities have been reducing temperature extremes, but it is not clear exactly how much of the noted recent change is due to man-made conditions as opposed to natural climatic variation.

A plot of seasonal decadal values of moderated temperature months for three central area stations (Fig. 18) reveals that the total (annual) increase with time, apparent in Figure 16, is largely due to recent increased frequencies in the summer and fall seasons. Little change with time is noted during spring with a decrease at some stations in the 1961-1970 decade. The winter frequencies are uniformly low with an increase in the last 10-year



Figure 18. Seasonal frequencies of moderated temperatures at Urbana, Winamac, Ind. , and Brunswick, Mo.

period at 2 stations. Seasonal frequencies for. cloudy skies in the Peoria area showed increases since the 1940's, but the greatest seasonal increases occurred in summer and fall when the moderated temperature increases were found to be greatest. The similarities in the seasonal frequencies between the four stations located along a 1000 km east-west line (Fig. 18) help reveal the regional nature of the temperature influence in the central area (Fig. 1). The seasonal results further support the conclusion that recent increases in cloudiness are largely responsible for the increase in moderated temperatures. Pollutant-related haze seems a less likely cause.

Precipitation Effects

This study did not have as an objective a search for contrail influences on precipitation. However, the decided downward trends in frequency of clear skies and increased cloudy skies, led to a comparison of precipitation with clear sky frequencies at two Illinois stations. The 10-year values of precipitation and the frequencies of clear days were derived for Peoria and Springfield, for 1901-1970, as well as the 7-year value (for 1971-77). This series of eight independent values at each station are plotted on Figure 19. This scattergram indicates a weak relationship between the frequency of clear days and annual precipitation and the correlation coefficient is -0.45. As expected, less precipitation occurs when more clear days occur. This finding suggests that the decrease in clear days found at these stations and elsewhere in the central areas of the Midwest since 1960, is related to a period of greater precipitation, and may reflect part of the cause of more precipitation.



Figure 19. Relationships between clear day frequencies and precipitation at Peoria and Springfield,

Conclusions

The sky cover frequencies and sunshine values indicate sizeable changes during the 1901-77 period. These occur in most midwestern areas and in most seasons. Basically, the frequency of clear days and the percent of possible sunshine have decreased substantially since 1930, although the sunshine has decreased less than the clear day conditions. Conversely, the cloudy day frequencies have increased but at a greater rate than the clear day frequencies. The partly cloudy frequencies have decreased slightly in the central areas in summer and fall.

The biggest shifts in sky cover and sunshine occurred across the North Central and South Central Areas. These also happened to occur in the periods of jet air traffic and the greatest urban-industrial development. The biggest shifts in sky cover and sunshine also occurred in the fall and summer seasons, when conditions normally are least cloudy. The sunshine changes are totally restricted to the central area and strongly point to regional man-made (not natural) influences. The sunshine decrease from 1920 to 1977 was 4 and 7% in summer and fall, respectively. Cloudy days revealed an anomalous shift from 1910 to 1977 with areal increases varying from 11 to 27% in the summer, and from 9 to 26% in the fall. Clear days and partly cloudy days exhibited lesser shifts, decreases of 10 to 20%, in summer and fall.

Recent upward trends in cloudiness (and decreases in sunshine) have been particularly notable in fall and summer and within the Midwest's central section (across the northern half of Missouri, Illinois, and Indiana, and the southern halves of Iowa, Wisconsin, and lower Michigan) were jet air traffic is greatest and also where urban-industrial development has been greatest in the Midwest.

It was hypothesized initially that the causes of marked shifts in midwestern cloudiness and sunshine could be attributed to any one of three factors. The first of these was natural causes, i.e., short or long term climatic changes on areal scales much larger than that in the study region. The second cause could be attributed to errors in observation or in the sunshine instrumentation. The third possible cause of trends includes inadvertent, man-made influences which could be subdivided into a pollutant-related decreases in visiblity, and radiation, pollutant-related influences on clouds, and a jet contrail influence on clouds.

The results collectively suggest that all of these causative factors were operative in the study region. For example, long-term trends were found in the frequency of cloudy days in all four areas. This suggests a natural large-scale shift in the climate, probably extending over large portions of North America during the 1901-77 period. However, such long-term trends were not found in the sunshine frequencies. Secondly, within the long-term trends in the sky cover conditions are two major anomalies, one of which is found in the sunshine data of certain areas and seasons.

A major anomaly noted was a sharp shift to more cloudiness during 1941-50. This occurred at a time when there was great urban-industrial expansion in the Midwest, and also when most observation stations were relocated from city sites to airport sites. Such a move to more rural sites should have improved sky visibility but occurred at a time when production of pollutants (as reflected in haze-smoke days) was rapidly increasing across the region.. For this and other reasons, the error-related explanation for the trends was discarded. Since the 1941-50 anomaly was most pronounced in the more heavily populated urban industrial parts of the Midwest, it appears that this shift may be due largely to the effects of man-made pollutants acting to

decrease visibility of clouds, to decrease incoming sunshine, and to assist in the formation of clouds. The marked increase in frequency of days with haze and smoke in the 1935-1950 period agreed very well with the shifts in sky cover, but not in sunshine. Thus, some of the change at this time was due to pollutants effect on visibility and possibly on low clouds (and not on high clouds).

The other notable anomaly found in the general long-term trends in sky cover and sunshine occurred principally in the central section. It was a decrease in clear skies and sunshine beginning about 1960, and reaching the lowest values since 1900 in the late 1970's. Importantly and conversely, in other areas to the north and south of the central Midwest, the clear day frequencies have increased since they achieved quite low values in the 1940's. This anomaly of the last 15 to 20 years also has occurred largely only in the fall and summer seasons, normally the seasons of clearer skies. There is much indirect evidence pointing towards contrail-related influences in this anomalous period and area. If true, the contrail influence has represented a 5-10% change in the sunshine, clear day, and cloudy day frequencies. Influence on sunshine is less than on cloudy conditions presumably because on many days with contrail cirrus the sunshine still penetrates and is recorded.

The three primary findings of this study are as follows. There has been 1) a long-term trend toward more cloudiness over the 1901-77 period (not as evident in sunshine), 2) a sharp increase in cloudiness during 1941-50, and 3) a second anomalous increase in cloudiness (and decrease in sunshine) in the central sections of the Midwest from 1961 to the end of the study period (1977). The first of these three findings is believed to be a part of a

general climatic trend with the last two apparently related to inadvertent man-made influences (pollutants beginning in 1940 and now contrails in more recent years).

The influence of these trends is reflected in surface temperature extremes (less extreme minimum and maximum monthly averages) which have generally increased with time. At weather stations in the central area of the Midwest, increases in moderated temperatures occurred in the 1940's with the anomalous increase in cloudy conditions believed to be related to the massive release of pollutants that began at that time. Finally, moderated temperatures occurrences have shown a sudden upswing in the same area since 1960, which relates well with the decreased sunshine and increased cloudiness noted in this period. In addition, the increased frequency of moderated temperatures since 1960 has occurred largely in the fall and summer, the same seasons when the cloud cover increases were noted. The results are internally very consistent.

These results alone, particularly those relating to the increased cloudiness since 1960, cannot be considered proof of a jet contrail influence. However, this anomalous change is regionally restricted to that area of the Midwest where jet traffic is the greatest, but this may still represent only a natural regional scale fluctuation of sky conditions. Other results of this project presented in later chapters help define the potential of contrail influences on a) cloudiness, b) sunshine received, and c) surface temperatures during this apparent anomalous period that began around 1960.

CHAPTER TWO

CLOUD TYPE CLIMATOLOGY OVER THE RESEARCH AREA

Introduction

The observational data available for the first order stations shown in Figure 20 were used to study the trend of cloud types during the 26-year period 1951-1976. The frequency of types were categorized by standard definition into the simple categories of low, middle, and high clouds according to standard observational definitions. The 3-hour observation times throughout the 24-hour day were used to obtain frequencies for the classes of clouds involved. A fourth class, the frequency of clear sky observations is also included since it may be a sensitive parameter for the determination of long-term trends in cloudiness.

The stations used for this limited study included Duluth, Minnesota; Sault Ste. Marie, Michigan; Madison, Wisconsin; Des Moines, Iowa; Chicago, Illinois; Peoria, Illinois; Springfield, Illinois; St. Louis, Missouri; Ft. Wayne Indiana; and Evansville, Indiana. These provided sufficient density to examine a north-south variation in cloudiness extending from Lake Superior to the Ohio River and a similar east-west transect from Des Moines to Ft. Wayne. In referring to Figure 2, the stations extend across a high density traffic area, i.e., an area of high potential contrail formation. The observed traffic density suggests the use of Duluth and Sault Ste. Marie as controls against which the trends of cloudiness at the central Midwestern stations can be compared.

In this chapter the temporal trend of individual cloud types will be examined culminating in two stratifications of high cloudiness in an attempt to identify an indicator of contrail activity. The parameter used is the



Figure 20. The Midwest area of study with the first-order weather stations used in the cloud-type frequency analysis.

slope of the trend line generated by the data over the 26-year period although further detailed analyses might suggest that shorter interval trend lines may illuminate certain features not readily apparent from the entire period.

Low Cumulus Cloud Frequency Trend

An example of the trend of low cumulus cloud frequency at Peoria, Illinois is shown in Figure 21. The correlation coefficient for the trend line was 0.63. The trend line intercepts the year 1951 at a frequency of 350 and extends upward to 400 at the end of 1976. These end point numbers are an indication of the general climatic variation of cloudiness over the broad area encompassed by this study and are shown for each of the stations in Table 7. The slope of low and middle clouds should represent the climatic trend, i.e., a control, not influenced by jet contrails. The lowest frequency in 1951 was observed at Sault Ste. Marie with a value of about 189 with the highest value of 432 at Springfield, Illinois. Three stations showed negative slopes, that is, decreasing cloudiness, and all others showed increases of low cumulus clouds over the 26-year interval.

The geographic variability of the trend line slope is shown in Figure 22. The pattern reveals the negative slopes for the trend of cloudiness at Springfield, Des Moines, and Duluth with high positive slopes eastward over most of Illinois, Indiana, and northern Michigan. The greater positive slopes are evident from Des Moines southeastward and from St. Louis northeastward merging over central Indiana.

The significance of the study of the low cloudiness trend is to qualitatively assess the potential effect on the observational quality of the high level clouds, including contrails. The high positive slopes of the low

Table 7. The trend line point values of annual cloudiness over the Midwest for 1951 and 1976.

		FTW	DSM	PIA	SSM	MLI	SPI	MDW	MSN	STL	EW	GRB	DLH
Low Strat	1951	1101	1013	838	1514	873	902	1100	1130	1008	1039	1140	[1436]
	1976	1557	1016	1265	2054	1067	1189	1276	1232	1023	1268	1141	[1411]
Low Cu	1951	336	[381]	343	189	408	[432]	305	310	320	371	337	[420]
	1976	344	[354]	443	293	517	[397]	314	352	426	402	354	[357]
Middle	1951	[756]	[925]	[840]	680	839	[978]	[774]	[923]	[846]	[807]	[860]	[1039]
	1976	[752]	[822]	[819]	760	903	[821]	[741]	[739]	[799]	[543]	[728]	[745]
High	1951	615	981	807	522	712	858	676	786	904	761	724	816
	1976	1027	1098	1060	874	1042	1044	935	1038	984	912	847	1038
£0.4 L&M	1951	474	770	652	432	566	675	512	626	697	628	567	620
	1976	768	816	819	649	788	797	690	785	702	678	641	728
£0.4 L&M <HC	1951	254	439	382	209	291	400	231	347	337	[340]	[333]	362
	1976	338	441	390	327	443	407	348	378	351	[335]	[299]	416
Clear	1951	[533]	[553]	[472]	[406]	[707]	[661]	[634]	[537]	[717]	[711]	[544]	336
	1976	[469]	[595]	[555]	[295]	[542]	[550]	[430]	[514]	[655]	[667]	[522]	415



Figure 21. The annual frequency of cumulus in the low cloud category at Peoria, Illinois for the 1951-1976 period.



Figure 22. The spatial pattern of the 26-year trend line slope for the "low cumulus" category of clouds.

cumulus category suggests that upper level clouds may be difficult to record due to intervening clouds over north-central and south-central Illinois, and central Indiana.

Low Stratus Cloud Frequency Trend

The low stratus category of clouds showed even a greater tendency than low cumulus toward increased frequency over the past 26 years. The largest change was evidenced at Sault Ste. Marie where the slope of the linear trend line through the 26-year data set was 21.6. However, all stations reported positive slopes with the exception of Duluth showing a small negative change in cloudiness with a slope of -0.69. The spatial variability of the low stratus category trend lines slope is shown in Figure 23.

A comparison of the trend plots for all the stations reveals that between the period 1958 and 1962 there was a relatively high increase in frequency of low stratus clouds at the more southern stations, i.e., in Illinois, Missouri, and Indiana. This period of relatively high frequency of low stratus clouds was followed by a period of values below the trend line for the years 1962 through 1968 at the same stations. However, the northern stations in Minnesota, Wisconsin, and Michigan showed opposite characteristics with frequency values below the trend line in the 1958-1962 period followed by a recovery to values exceeding the trend line during the later period. These variations are certainly a reflection of macroscale synoptic weather patterns which could dominate these two classifications of low clouds.

Middle Cloud Frequency Trend

The trend line for middle clouds at 10 out of 12 stations showed a decreasing frequency of occurrence. As shown in Table 7, this cloud



Figure 23. The spatial pattern of the 26-year trend line slopes for the "low stratus" category of clouds.

stratification is anomalous among those used in this study. The obscuration of middle-level clouds by low clouds might explain an apparent decrease, and, thus, predict a similar decrease in high cloud frequencies. If the downward trend, however, was due to macroscale conditions of the middle atmosphere, then the high cloud observations may still be representative of a true frequency change. As will be shown later, the high cloud stratification indicates an upward trend at nearly all stations suggesting the latter explanation is feasible.

The greatest decrease was observed at Duluth (Fig. 24) with a nearly equal trend at Evansville (Fig. 26) during this 26-year record. However, it should be noted that while the decrease is nearly the same at these two stations, the 1976 end point of the trend at Duluth nearly coincides with the 1951 initial point of the Evansville trend. This illustrates that there may be considerably more middle cloudiness in the northern latitudes (represented by Duluth), i.e., a climatological feature of Midwest cloudiness during 1951-1976.

For the most part through the Illinois-Indiana area, the trend of, or the difference between the 1951 and 1976 middle cloud frequencies is somewhat mixed and ill-defined. The data shown in Figure 27 represent the simple difference between the 1951 and 1976 frequency values. As shown in Figure 27, the greatest decrease of middle clouds largely occurred at the western edge of the study area. Smaller changes of cloud frequency occur in approximately the same area as maximum trend line slope of low cumulus shown in Figure 22. Values over the two-state area ranged from -157 (Springfield) to +64 (Moline) which along with Sault Ste. Marie, showed an increase of middle cloudiness (Fig. 24). One might suspect that Sault Ste. Marie may suffer from the effects of Lake Superior to the west, but on the other hand the absolute value



Fiqure 24. The annual frequency of middle clouds during 1951-1976 at Duluth, Sault Ste. Marie, Green Bay, and Madison.